

The Nonlinear Maslov index and the Calabi homomorphism

Gabi Ben Simon ^{*†}

School of Mathematical Sciences
Tel Aviv University 69978 Tel Aviv, Israel
gabiben@post.tau.ac.il

February 2, 2008

Abstract

In this paper we find that the asymptotic nonlinear Maslov index defined on the universal cover of the group of all contact Hamiltonian diffeomorphisms of the standard $(2n - 1)$ -dimensional contact sphere is a quasimorphism. Then we show our main result: Let M be standard $(n - 1)$ -dimensional complex projective space. We prove that the value of the pullback of the asymptotic nonlinear Maslov index to the universal cover of the group of Hamiltonian diffeomorphisms of M , when evaluated on a diffeomorphism supported in a sufficiently small open subset of M , equals $\frac{-n}{2\pi \text{vol}(M)}$ times the Calabi invariant of this diffeomorphism.

1 Introduction and results

1.1 Basics on the nonlinear Maslov index

Let (S^{2n-1}, ξ) be the $(2n - 1)$ -dimensional sphere endowed with the standard contact structure. We denote by $\mathcal{C}$ the identity component of the group of those contact Hamiltonian diffeomorphisms of (S^{2n-1}, ξ) , which commute with the inversion $x \mapsto -x$ on S^{2n-1} (see [7] for preliminaries on contact manifolds). Let $\tilde{\mathcal{C}}$ be the universal cover of $\mathcal{C}$. Recall that an element $\tilde{f}$ of $\tilde{\mathcal{C}}$ can be identified as the time-1 map of the flow generated by a contact Hamiltonian, up to homotopy with fixed endpoints. Multiplication on $\tilde{\mathcal{C}}$ is defined as follows: let $\tilde{f}, \tilde{g}$ in $\tilde{\mathcal{C}}$ be the time-1 map of the

^{*}This paper is a part of the author's Ph.D. thesis, being carried out under the supervision of Professor Leonid Polterovich, at Tel-Aviv University.

[†]Partially supported by Israel Science Foundation grant 11/03.

flows $\{\tilde{f}_t\}_{0 \leq t \leq 1}$ and $\{\tilde{g}_t\}_{0 \leq t \leq 1}$ respectively. Then the multiplication of $\tilde{f}$ and $\tilde{g}$ is the homotopy class of the path $\{\tilde{f}_t \tilde{g}_t\}_{0 \leq t \leq 1}$.

Moreover, $\mathcal{C}$ carries the structure of an infinite dimensional manifold. Following Givental (see [5]), we denote by $\Delta(\mathcal{C})$ the *discriminant* of $\mathcal{C}$, that is the hypersurface of $\mathcal{C}$ formed by all contactomorphisms which have a fixed point, and preserve the standard contact form at that point. Using generating functions, Givental has defined a co-orientation of $\Delta(\mathcal{C})$ in $\mathcal{C}$.

Given a path $\{\tilde{f}_t\}_{t \geq 0}$ in $\mathcal{C}$, an intersection index can be assigned to it by counting all signed intersections of the path with a subspace of $\Delta(\mathcal{C})$, called the *train* and denoted by $\Gamma(\mathcal{C})$, relative to the co-orientation at nonsingular points of $\Delta(\mathcal{C})$, (see again [5]).

We call the intersection index of $\{\tilde{f}_s\}_{s \in [0, t]}$ with $\Gamma(\mathcal{C})$ the *nonlinear Maslov index* of $\{\tilde{f}_s\}_{s \in [0, t]}$ and denote it by $\mu(t)$. Following Givental we now define the *asymptotic nonlinear Maslov index*: let $\tilde{f}$ be an element of $\tilde{\mathcal{C}}$ generated by a time-1 periodic contact Hamiltonian $\tilde{F} : S^{2n-1} \times S^1 \rightarrow \mathbb{R}$. Let $\{\tilde{f}_t\}_{t \geq 0}$ be the flow generated by $\tilde{F}$, which is a path in $\mathcal{C}$. The asymptotic nonlinear Maslov index of $\tilde{f}$ is then given by

$$m(\tilde{f}) := \lim_{t \rightarrow \infty} \frac{\mu(t)}{t}. \quad (1)$$

1.2 The quasimorphism property

Definition 1.1. Let G be a group. A *quasimorphism* r on G is a function $r : G \rightarrow \mathbb{R}$ which satisfies the homomorphism equation up to a bounded error: there exists $R > 0$ such that

$$|r(fg) - r(f) - r(g)| \leq R$$

for all $f, g \in G$.

See [2] for preliminaries on quasimorphisms. A quasimorphism r_h is called *homogeneous* if $r_h(g^m) = m r_h(g)$ for all $g \in G$ and $m \in \mathbb{Z}$. Every quasimorphism r gives rise to a homogeneous one

$$r_h(g) = \lim_{m \rightarrow \infty} \frac{r(g^m)}{m}.$$

We now can state the first theorem of this paper which is actually an easy consequence of [5].

Theorem 1.2. *The asymptotic nonlinear Maslov index $m : \tilde{\mathcal{C}} \rightarrow \mathbb{R}$ is a quasimorphism.*

The proof will be given in Section 2 after some preliminaries on the asymptotic nonlinear Maslov index.

1.3 The pullback of m

Let $(\mathbb{C}P^{n-1}, \omega)$ be $(n-1)$ -dimensional complex projective space endowed with the Fubini-Study form, normalized so that $\int_{\mathbb{C}P^1} \omega = \pi$. Let $G = \text{Ham}(\mathbb{C}P^{n-1}, \omega)$ be the identity component of the group of all Hamiltonian diffeomorphisms of $(\mathbb{C}P^{n-1}, \omega)$ and $\tilde{G}$ its universal cover (see [7] and [8] for preliminaries on G). Let $\{f_t\}_{t \in [0,1]}$ be a path in G representing an element of $\tilde{G}$. There is a unique Hamiltonian $F : \mathbb{C}P^{n-1} \times S^1 \rightarrow \mathbb{R}$, which generates $\{f_t\}_{t \in [0,1]}$ and satisfies the normalization condition:

$$\int_{\mathbb{C}P^{n-1}} F_t \omega^{n-1} = 0$$

for all $t \in S^1$.

We denote the space of all normalized 1-periodic Hamiltonians by $\mathbf{F}$. In order to pull back the asymptotic nonlinear Maslov index from $\tilde{\mathcal{C}}$ to $\tilde{G}$ we will use the Hopf bundle

$$p : S^{2n-1} \rightarrow \mathbb{C}P^{n-1}.$$

Let $f \in \tilde{G}$, and let $\{f_t\}_{t \in [0,1]}$ be a representative of f . Let $F \in \mathbf{F}$ be the unique normalized Hamiltonian generating this path. Then we can lift F to S^{2n-1} to a time dependent function,

$$\tilde{F}_t := p^* F_t.$$

$\tilde{F}$ is constant along the fibers of the fiber bundle $p : S^{2n-1} \rightarrow \mathbb{C}P^{n-1}$. This $\tilde{F}$ is a contact Hamiltonian on the contact sphere. The contact flow $\{\tilde{f}_t\}_{t \in [0,1]}$ generated by this contact Hamiltonian constitutes a lift of the flow $\{f_t\}_{t \in [0,1]}$, and represents an element $\tilde{f}$ of $\tilde{\mathcal{C}}$. The value of the pullback of the nonlinear Maslov index at f is then set to be $m(\tilde{f})$. We will denote by

$$i : \tilde{G} \rightarrow \tilde{\mathcal{C}} \tag{2}$$

the above injection, and by

$$i^* m : \tilde{G} \rightarrow \mathbb{R} \tag{3}$$

the pullback of the asymptotic nonlinear Maslov index. As a corollary of Theorem 1.2 we have

Corollary 1.3. *The pullback of the asymptotic nonlinear Maslov index*

$$i^* m : \tilde{G} \rightarrow \mathbb{R}$$

is a quasimorphism.

Remark. Any Hamiltonian (not necessarily normalized) generating an element of $\tilde{G}$ can be lifted via p in the way described above. The normalization condition is required since we want to embed $\tilde{G}$ in $\tilde{\mathcal{C}}$.

We close this subsection with a simple, but important example.

Example 1.4. Consider the flow $f_t = \mathbf{1}$ for all t where $\mathbf{1}$ is the identity map of $\mathbb{C}P^{n-1}$. Any function of the form $F(x, t) = F(t)$ (clearly, not necessarily normalized) generates this flow. Let p^*F_t be its lift to the sphere. An easy computation shows that the flow $\{\tilde{f}_t\}$, generated by p^*F_t , leaves the fibers of the Hopf bundle invariant, and rotates them by angle $e^{iF(t)}$ for time t .

1.4 The Calabi property of i^*m

In this subsection we will assume that $G = \text{Ham}(M, \omega)$, where M^{2n} is any closed connected symplectic manifold.

The group G has a natural class of subgroups G_U associated to nonempty open subsets $U \subset M$. An element f of G lies in G_U if it is the time-1 map of a Hamiltonian flow $\{f_t\}_{t \in [0,1]}$ with $f_0 = \mathbf{1}$, generated by a Hamiltonian $F_t : M \rightarrow \mathbb{R}$, $t \in [0, 1]$, such that

$$\text{support}(F_t) \subset U \text{ for all } t. \quad (4)$$

Consider the map $\text{Cal}_U : G_U \rightarrow \mathbb{R}$ given by

$$f \mapsto \int_0^1 dt \int_M F_t \omega^n. \quad (5)$$

When the symplectic form ω is exact on U , this map is well-defined, meaning that it does not depend on the specific choice of the Hamiltonian F generating f . In fact, Cal_U is a homomorphism called *the Calabi homomorphism* (see [1], [3] and [10] for further information).

The above definitions can be “lifted” to $\tilde{G}$: for a nonempty subset $U \subset M$, consider the subgroup $\tilde{G}_U \subset \tilde{G}$ defined as follows. An element of $\tilde{G}$ lies in $\tilde{G}_U$ if it can be represented by a Hamiltonian flow $\{f_t\}_{t \in [0,1]}$ with $f_0 = \mathbf{1}$, generated by a Hamiltonian F_t satisfying condition (4). Formula (5) gives rise to a well-defined homomorphism

$$\widetilde{\text{Cal}}_U : \tilde{G}_U \rightarrow \mathbb{R}.$$

In what follows we deal with the class $\mathcal{D}$ of all nonempty open subsets U which can be *displaced* by a Hamiltonian diffeomorphism:

$$hU \cap \text{Closure}(U) = \emptyset \text{ for some } h \in G. \quad (6)$$

We need one more definition before stating our main result.

Definition 1.5. A quasimorphism on $\tilde{G}$ coinciding with the Calabi homomorphism $\{\widetilde{\text{Cal}}_U : \tilde{G}_U \rightarrow \mathbb{R}\}$ on each $U \in \mathcal{D}$ is said to have the *Calabi property*.

From now on $M = \mathbb{C}P^{n-1}$. Here is our main result:

Theorem 1.6. *The pullback of the asymptotic nonlinear Maslov index $i^*m : \tilde{G} \longrightarrow \mathbb{R}$ has the Calabi property up to multiplication by $-\frac{n}{2\pi\text{vol}(M)}$. That is*

$$i^*m([f]) = -\frac{n}{2\pi\text{vol}(M)} \widetilde{\text{Cal}}_U([f])$$

for every element $[f] \in \tilde{G}_U$.

The proof will be given in section 3.

Remark. Given a Hamiltonian $H : M \times S^1 \longrightarrow \mathbb{R}$ denote by f^H the element of $\tilde{G}$ generated by H . A function $r : \tilde{G} \longrightarrow \mathbb{R}$ is called *continuous*, if it satisfies the following condition: if a sequence $\{H_i\}$ of smooth Hamiltonians $H_i : M \times S^1 \longrightarrow \mathbb{R}$ C^0 -converges to a smooth function $H : M \times S^1 \longrightarrow \mathbb{R}$ then

$$r(f^{H_i}) \longrightarrow r(f^H) \text{ as } i \rightarrow \infty. \quad (7)$$

It follows from Corollary 3 of Section 9 of [5] that the pullback of the asymptotic nonlinear Maslov index is continuous. Summarizing, the pullback of the asymptotic nonlinear Maslov index $i^*m : \tilde{G} \longrightarrow \mathbb{R}$ is a continuous quasimorphism with the Calabi property (up to multiplication by a constant).

1.5 Other quasimorphisms with the Calabi property

Another construction of a continuous quasimorphism on $\tilde{G}$ with the Calabi property has been carried out by Entov and Polterovich in [4]. In particular it applies to the manifold $(\mathbb{C}P^{n-1}, \omega)$.

Their construction uses the spectral invariants of elements of $\tilde{G}$. These spectral invariants are defined via the Floer theory on the symplectic manifold. The first step in our proof of theorem 1.6 goes along the lines of [4]. Moreover, step 3 of the proof of 1.6 was partly motivated by the inspection of the behavior of spectral invariants defined in terms of generating functions (see [4] for more details).

2 Preliminaries on the nonlinear Maslov index

In this short section we state the main properties of the asymptotic nonlinear Maslov index and the nonlinear Maslov index μ , which we use in the proof of the main result (theorem 1.6).

First, we remind the reader that the sign of an intersection of a path in $\tilde{\mathcal{C}}$ with the train (which is a subspace of the discriminant) is determined via the co-orientation of the train, which is defined by methods of generating functions. The two basic facts that we need are grouped together in the following theorem, which is due to Givental (see [5], [6], and [9]).

Theorem 2.1. (a) *The nonlinear Maslov index (relative to the train) of any two paths in $\mathcal{C}$ is the same, provided that there is a homotopy between them, such that the end points of the paths in this homotopy lie outside the discriminant.*

(b) *Let f be a Hamiltonian map of $\mathbb{C}P^{n-1}$ and let $\tilde{f}$ be its lift to $\mathcal{C}$. Assume that it has no fixed points. Then the nonlinear Maslov index of the loop $\{e^{\pm 2\pi i t} \tilde{f}\}_{t \in [0,1]}$ (relative to the train) is $\pm n$.*

2.1 Proof of Theorem 1.2

Let $\tilde{f} \in \tilde{\mathcal{C}}$. By definition, it is generated by a 1-periodic contact Hamiltonian. Let $\{\tilde{f}_t\}$ be the contact flow generated by this Hamiltonian. Then we have $\tilde{f}^m = \tilde{f}_m$ for all integer m . This means that

$$m(\tilde{f}) = \lim_{m \rightarrow \infty} \frac{\mu(\tilde{f}^m)}{m}. \quad (8)$$

It suffices to prove that μ is a quasimorphism on $\tilde{\mathcal{C}}$ (see section 1.2). For this, let $[\tilde{f}]$ and $[\tilde{h}]$ be elements of $\tilde{\mathcal{C}}$, represented by paths $\{\tilde{f}_t\}_{t \in [0,1]}$ and $\{\tilde{h}_t\}_{t \in [0,1]}$, respectively. Consider the following path γ , which represents $[\tilde{f}][\tilde{h}]$. It is the concatenation

$$\{\tilde{f}_{2t}\}_{0 \leq t \leq \frac{1}{2}} * \{\tilde{f}_1 \tilde{h}_{2t-1}\}_{\frac{1}{2} \leq t \leq 1} := \begin{cases} \tilde{f}_{2t} & 0 \leq t \leq \frac{1}{2} \\ \tilde{f}_1 \tilde{h}_{2t-1} & \frac{1}{2} \leq t \leq 1 \end{cases}.$$

It follows from Givental's construction that μ is additive with respect to concatenation. So we have that

$$\mu(\gamma) = \mu([\{\tilde{f}_{2t}\}]) + \mu([\{\tilde{f}_1 \tilde{h}_{2t-1}\}]). \quad (9)$$

Moreover it is also follows from his construction that

$$|\mu([\{\tilde{h}_{2t-1}\}]) - \mu([\{\tilde{f}_1 \tilde{h}_{2t-1}\}])| \leq 2n. \quad (10)$$

Combining (9) and (10) we get

$$|\mu(\gamma) - \mu([\{\tilde{f}_{2t}\}]) - \mu([\{\tilde{h}_{2t-1}\}])| \leq 2n.$$

Since the paths $\{\tilde{f}_{2t}\}$ and $\{\tilde{h}_{2t-1}\}$ represent respectively $[\tilde{f}]$ and $[\tilde{h}]$, we have:

$$|\mu([\tilde{f}][\tilde{h}]) - \mu([\tilde{f}]) - \mu([\tilde{h}])| \leq 2n,$$

which is the desired. $\square$

3 Proof of Theorem 1.6

3.1 Step 1

Let $\{f_t\}_{t \in [0,1]}$ be a representative of an element $[f] \in \widetilde{G}_U$ for some open set U in $\mathcal{D}$, generated by a time-1 periodic Hamiltonian F_t (see 1.4). We have to show that:

$$-\frac{n}{2\pi \text{vol}(M)} \widetilde{\text{Cal}}_U([f]) = i^* m([f]) . \quad (11)$$

Recall that the right hand side is actually the asymptotic nonlinear Maslov index of the lift $\{\tilde{f}_t\}_{t \geq 0}$ of the flow $\{f_t\}_{t \geq 0}$ (see 1.1 and 1.3). Let h be a Hamiltonian diffeomorphism displacing U , and suppose that it is the time-1 map of a Hamiltonian isotopy $\{h_t\}_{t \in [0,1]}$. As a first step, we claim that it is enough to show that the asymptotic nonlinear Maslov index of the path $\{\widetilde{hf}_t\}_{t \geq 0}$, which is the lift of the path $\{hf_t\}_{t \geq 0}$, equals the left-hand side of (11). This follows easily from inequality (10). Since the difference of the nonlinear Maslov indexes of these paths is bounded by $2n$, they have the same asymptotic nonlinear Maslov index.

From now on we will focus on the concatenation

$$\{h_t\}_{t \in [0,1]} * \{hf_{t-1}\}_{t \geq 1} := \gamma_t ,$$

and its lift to $\mathcal{C}$

$$\{\tilde{h}_t\}_{t \in [0,1]} * \{\widetilde{hf}_{t-1}\}_{t \geq 1} := \tilde{\gamma}_t .$$

Let us emphasize, that the paths $\{\gamma_t\}_{0 \leq t \leq t_0}$ and $\{\tilde{\gamma}_t\}_{0 \leq t \leq t_0}$ (for $1 \leq t_0$) represent respectively hf_{t_0-1} and $\widetilde{hf}_{t_0-1}$ as elements of $\widetilde{G}$ and $\widetilde{\mathcal{C}}$ respectively.

Secondly, note that because h displaces U , the set of fixed points of $\{hf_t\}_{t \geq 0}$ coincides with the set of fixed points of h for all $t \geq 0$. Let x_0 be a fixed point of h , and let $\tilde{x}_0 \in p^{-1}(x_0)$ be a point of the fiber $p^{-1}(x_0)$ above x_0 . Of course $\tilde{\gamma}_t(\tilde{x}_0)$ is a lift of $\gamma_t(x_0)$. In particular for $0 \leq t \leq 1$ the path $\tilde{\gamma}_t(\tilde{x}_0)$ is a lift of the loop $\gamma_t(x_0)$, but this path need not be closed. Nevertheless the point $\tilde{x}_0$ will return to its fiber for $t = 1$. For $t \geq 1$ we have $\gamma_t(x_0) = x_0$, so the fiber $p^{-1}(x_0)$ is invariant under the flow $\tilde{\gamma}_t$. Therefore we have:

$$\tilde{\gamma}_t(\tilde{x}_0) = e^{i\theta(t)} \tilde{x}_0 , t \geq 1 \quad (12)$$

where $\theta(t)$ is determined by the contact flow generated by the lift of the Hamiltonian generating the path γ_t .

3.2 Step 2

Now, assume that for $t = s$, $s > 1$ the path $\tilde{\gamma}_t$ intersects the train $\Gamma(\mathcal{C})$. This means that there exists $x_0 \in \{\text{fix}(h)\}$ such that in equation (12) we have $\theta(s) = 2\pi k$ for some $k \in \mathbb{Z}$. In the second step of the proof we first calculate the function $\theta(s)$, and

then modify the path $\tilde{\gamma}_t$ to one for which the function $\theta(s)$ is linear. This will help us to make the final reduction step in the next section, where we actually calculate the asymptotic nonlinear Maslov index.

Notation: From now on we denote $C = -\frac{\widetilde{\text{Cal}}_U([f])}{\text{vol}(M)}$.

Let $\{K_t\}$ be the normalized Hamiltonian generating the flow $\{\gamma_t\}$, and let $\{\tilde{K}_t\}$ be its lift. Clearly for $t \geq 1$ we have:

$$K_t = F_t + c(t)$$

where F_t is a 1-periodic Hamiltonian generating the flow $\{f_t\}$, and

$$c(t) = -\frac{1}{\text{vol } M} \int_M F_t \omega^{n-1}.$$

Moreover, for all $x \in M \setminus U$ we have $K_t(x) = c(t)$.

For $s \geq 1$ define

$$\theta(s) := \int_1^s c(t) dt = - \int_1^s dt \frac{1}{\text{vol } M} \int_M F_t \omega^{n-1}.$$

Since F_t is periodic we have for every $m \in \mathbb{N}$ and $0 \leq v \leq 1$

$$\theta(m+v) = C(m-1) - \int_0^v dt \frac{1}{\text{vol } M} \int_M F_t \omega^{n-1}.$$

Note also that the function $\varphi(s) := \theta(s) - C(s-1)$ is 1-periodic and satisfies $\varphi(m) = 0$ for every $1 \leq m \in \mathbb{N}$. In particular for all such m we have

$$C(m-1) = \theta(m). \tag{13}$$

The notation $\theta(s)$ here is deliberate. Actually, it is the solution of (12) for all fixed points of h . Note that $\theta(s)$ measures the total displacement of the fiber after time s . From now on we assume, without loss of generality, that h has a finite number of fixed points. This always can be achieved by a small perturbation of h .

Now let $\{x_1, \dots, x_l\}$ be the set of fixed points of h and let $\tilde{x}_j$ be a point above x_j for all $1 \leq j \leq l$. Let α_j , $0 \leq \alpha_j < 2\pi$, be the angle defined by the equation $\tilde{\gamma}_1(\tilde{x}_j) = e^{i\alpha_j} \tilde{x}_j$. We have for $s \geq 1$

$$\tilde{\gamma}_s(\tilde{x}_j) = e^{iC(s-1)} e^{i\varphi(s)} e^{i\alpha_j}(\tilde{x}_j). \tag{14}$$

From the properties of $\varphi(s)$ we get that the path $\{e^{i\varphi(s_0+t)}\}_{t \in [0,k]}$, $k \in \mathbb{N}$ is homotopic to the constant loop $\{e^{i\varphi(s_0)}\}$, for every $s_0 \geq 0$. We emphasize that the homotopy has $\{e^{i\varphi(s_0)}\}$ as the starting and end point of all of its paths.

Let us denote such a homotopy by $\{e^{iF_s(t)}\}_{(s,t) \in [0,1] \times [0,k]}$. So we have

$$F_0(t) = \varphi(s_0 + t), \quad F_1(t) = \varphi(s_0)$$

and

$$F_s(0) = F_s(k) = \varphi(s_0).$$

For reasons that will become clear we are looking for s_0 that satisfies the following condition:

$$C(s_0 + k - 1) + \varphi(s_0 + k) + \alpha_j \notin 2\pi\mathbb{Z} \quad (15)$$

for all $1 \leq j \leq l$ and $k \in \mathbb{N}$.

Such s_0 clearly exists. To see this, first note that the set $2\pi\mathbb{Z}$ is countable as well as $\mathbb{N}$, the set $\{\alpha_1, \dots, \alpha_l\}$ is finite, and finally $\varphi(s_0 + k) = \varphi(s_0)$ for all $k \in \mathbb{N}$. So we have to find a point in the image of the function $C(s - 1) + \varphi(s)$ which belongs to the complement of the countable set $\bigcup_{j,k} \{2\pi\mathbb{Z} - \alpha_j - kC\}$. This function is continuous and unbounded, thus its image clearly contains a segment, and consequently a point in the complement of the above set.

For such s_0 we have for all j, k

$$\tilde{\gamma}_{s_0+k}(\tilde{x}_j) = e^{iC(s_0+k-1)} e^{i\varphi(s_0)} e^{i\alpha_j}(\tilde{x}_j) \neq \tilde{x}_j. \quad (16)$$

That is for all $k \in \mathbb{N}$ $\tilde{\gamma}_{s_0+k}$ has no fixed points (and thus it is outside of the train). Now we have come to the key idea of step 2.

Lemma 3.1. *The family of maps*

$$\psi(s, t) = \{\tilde{\gamma}_{s_0+t} e^{-iF_s(t)} e^{i\varphi(s_0)}\}_{(s,t) \in [0,1] \times [0,k]}$$

is a homotopy of paths in $\mathcal{C}$ which has the following properties:

(a) *For $s = 1$ it is the path $\{\tilde{\gamma}_{s_0+t}\}_{t \in [0,k]}$.*

(b) *The starting point and the end point of each one of the paths comprising this homotopy has no fixed points. That is, the maps $\psi(s, 0)$, $\psi(s, k)$ have no fixed points.*

Proof. For $s = 1$ we have $F_1(t) = \varphi(s_0)$. Thus by the very definition of ψ we get part (a). For $t = 0$ and $t = k$ we know that for every $0 \leq s \leq k$ $F_s(0) = F_s(k) = \varphi(s_0)$. Thus by the definition of ψ and formula (16) we get part (b) of the lemma. $\square$

It follows from the invariance of the nonlinear Maslov index with respect to homotopies (see theorem 2.1) that the path $\psi(0, t)$ has the same index as the path $\psi(1, t)$ (which is $\{\tilde{\gamma}_{s_0+t}\}_{t \in [0,k]}$). Now, restricting the path $\psi(0, t)$ to the fibers over the fixed points we have the following equation

$$\psi(0, t)(\tilde{x}_j) = e^{iC(s_0+t-1)} e^{i\varphi(s_0)} e^{i\alpha_j} \tilde{x}_j. \quad (17)$$

We get the following equation by restricting the path $e^{i[C-\varphi(s_0)]}\psi(0, t)$ to the fibers over the fixed points:

$$e^{i[C-\varphi(s_0)]}\psi(0, t)(\tilde{x}_j) = e^{iC(s_0+t)}e^{i\alpha_j}(\tilde{x}_j)$$

Moreover, we see from equation (10) that the difference between the nonlinear Maslov indexes of $e^{i[C-\varphi(s_0)]}\psi(0, t)$ and $\psi(0, t)$ is bounded by $2n$ for all $k \in \mathbb{N}$. Thus, by the same argument we have used in step 1, we obtain that the asymptotic nonlinear Maslov indexes of the paths $e^{i[C-\varphi(s_0)]}\psi(0, t)$ and $\psi(0, t)$, $t \geq 0$, are equal.

It follows from the above discussion that without loss of generality we can assume that for $s \geq s_0$, $s_0 \geq 1$

$$\tilde{\gamma}_s(\tilde{x}_j) = e^{i[C(s-s_0)+\theta_j]}\tilde{x}_j, \quad (18)$$

where θ_j is the initial condition, which depends on s_0 and x_j , satisfies $0 \leq \theta_j < 2\pi$ and is defined by the equation

$$\tilde{\gamma}_{s_0}(\tilde{x}_j) = e^{i\theta_j}\tilde{x}_j.$$

3.3 Step 3

Before introducing the last step of the proof let us motivate it. We have seen that we may assume that the fibers above the fixed points of h rotate, under the action of $\{\tilde{\gamma}_t\}$, at the same angular speed and in the same direction of rotation (where the orientation is induced by the action of the group $\{e^{it}\}$). So after a period of time of length $\frac{2\pi}{|C|}$ each of the above fibers makes full rotation. This might lead us to believe that the number of signed intersections of $\{\tilde{\gamma}_t\}$, where t runs from some t_0 to $t_0 + \frac{2\pi}{|C|}$, with the train, is independent of t_0 , provided that $\tilde{\gamma}_{t_0}$ lies outside the train. In order to show that this is indeed the situation, we need the following lemma.

Lemma 3.2. *Let $\{\tilde{\gamma}_t\}_{t \geq 1}$ be the path in $\mathcal{C}$ as defined at the beginning of this section, and assume that it satisfies (18). Choose s_0 such that $\tilde{\gamma}_{s_0}$ is without fixed points. Then the family of maps*

$$\{e^{Ci(1-s)t}\tilde{\gamma}_{s_0+st}\}_{(s,t) \in [0,1] \times [0, \frac{2\pi}{|C|}]}$$

is a homotopy between the paths $\{\tilde{\gamma}_{s_0+t}\}_{t \in [0, \frac{2\pi}{|C|}]}$ and $\{e^{Cit}\tilde{\gamma}_{s_0}\}_{t \in [0, \frac{2\pi}{|C|}]}$ such that the starting and the end points of the paths of this homotopy stay outside the discriminant.

Proof. The first part of the lemma is trivial, since all we need to do is to substitute $s = 0$ and $s = 1$ to get the above paths. It is also obvious that the homotopy stays in $\mathcal{C}$. What we must show is that the constant path

$$\Lambda_1^s := \{\tilde{\gamma}_{s_0}\}$$

and the path

$$\Lambda_2^s := \{e^{Ci(1-s)\frac{2\pi}{|C|}}\tilde{\gamma}_{s_0+\frac{2\pi s}{|C|}}\}_{s \in [0,1]}$$

have no fixed points. We need only check this for the path Λ_2 . To this end let $x_0 \in \text{fix}(h)$, and let $\tilde{x}_0$ be any point above x_0 . Let $0 < \theta_0 < 2\pi$ be the angle of rotation of $\tilde{x}_0$ at time s_0 , where s_0 is chosen so as to satisfy (15). According to (18) we have

$$\tilde{\gamma}_{s_0 + \frac{2\pi s}{|C|}}(\tilde{x}_0) = e^{i(\text{sign}(C)2\pi s + \theta_0)}(\tilde{x}_0).$$

From this we get

$$\Lambda_2^s(\tilde{x}_0) = e^{i[\text{sign}(C)(1-s)2\pi + \text{sign}(C)2\pi s + \theta_0]}(\tilde{x}_0),$$

or simply:

$$\Lambda_2^s(\tilde{x}_0) = e^{i(\text{sign}(C)2\pi + \theta_0)}(\tilde{x}_0).$$

According to our choice of θ_0 clearly $\Lambda_2^s(\tilde{x}_0) \neq \tilde{x}_0$ for all $s \in [0, 1]$. This concludes the proof of our lemma. $\square$

Here we appeal to part (a) of theorem 2.1 and to lemma 3.2 in order to conclude that the paths

$$\{\tilde{\gamma}_{s_0+t}\}_{t \in [0, \frac{2\pi}{|C|}]}$$

and

$$\{e^{Cit}\tilde{\gamma}_{s_0}\}_{t \in [0, \frac{2\pi}{|C|}]}$$

has the same nonlinear Maslov index relative to the train. The calculation of the nonlinear Maslov index of $\{e^{Cit}\tilde{\gamma}_{s_0}\}_{t \in [0, \frac{2\pi}{|C|}]}$ uses part (b) of theorem 2.1, from which we get

Corollary 3.3. *The nonlinear Maslov index of the path $\{\tilde{\gamma}_{s_0+t}\}_{t \in [0, \frac{2\pi}{|C|}]}$ relative to the train is $\text{sign}(C)n$.*

Now we can calculate the asymptotic nonlinear Maslov index of $[f]$. Choose $s_0 \geq 1$ such that $\tilde{\gamma}_{s_0}$ has no fixed points. Note that $\tilde{\gamma}_{s_0 + \frac{2\pi k}{|C|}}$ also has no fixed points, and thus is outside the train, for all $k \in \mathbb{N}$. Define

$$\begin{aligned} a_k &:= \mu\left(\frac{2\pi k}{|C|} + s_0\right) - \mu(s_0) \\ &= \sum_{d=1}^k \left[\mu\left(\frac{2\pi d}{|C|} + s_0\right) - \mu\left(\frac{2\pi(d-1)}{|C|} + s_0\right) \right]. \end{aligned}$$

According to corollary 3.3 this equals

$$kn \text{sign}(C).$$

Thus

$$\begin{aligned} i^*m([f]) &= \lim_{t \rightarrow \infty} \frac{\mu(t)}{t} = \lim_{k \rightarrow \infty} \frac{a_k}{\frac{2\pi k}{|C|}} \\ &= \frac{n \text{sign}(C)|C|}{2\pi} = \frac{nC}{2\pi} \\ &= -\frac{n}{2\pi \text{vol}(M)} \widetilde{\text{Cal}}_U([f]), \end{aligned}$$

as claimed.

Finally, the case $C = 0$ follows from the considerations of step 2. According to what we have seen, we can take a homotopy of the path $\tilde{\gamma}_{s_0+t}$ such that there occurs no rotation in the fibers (see (17)), so clearly the asymptotic nonlinear Maslov index of $[f]$ is zero.

Acknowledgments. My sincere gratitude goes to Leonid Polterovich for all his support, ideas, and encouragement. I want to thank Michael Entov for reading drafts of this paper, and for coming to my talk about this work at Tel Aviv university and sharing his opinions with me. I would also like to thank Yaron Ostrover and Frol Zapolsky for reading the last draft of this paper and for their remarks.

References

- [1] Banyaga, A., *Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique*, comm. Math. Helv. **53**:2 (1978), 174-227.
- [2] Bavard, C. *Longueur stable des commutateurs*, L'Enseign. Math. **37**:1-2 (1991), 109-150.
- [3] Calabi, E., *On the group of automorphisms of a symplectic manifold*, in *problems in analysis*, 1-26. Princeton Univ. Press, 1970.
- [4] Entov, M.; Polterovich, L. *Calabi quasimorphism and quantum homology*. Int.math.Res.Not. **2003** no.30, 1635-1676.
- [5] Givental A.B. *Nonlinear generalization of the Maslov index*. Theory of singularities and its applications, Adv. Soviet Math., vol.1, American Mathematical society, Rhode Island, 1990, p.71-103.
- [6] Givental A.B. *The nonlinear Maslov index*. Geometry of low dimensional manifolds, 2 (Durham, 1989), 35-43. London Math. Soc. Lecture Note Ser., 151, Cambridge university press 1990.
- [7] McDuff, D., Salamon, D., *Introduction to symplectic topology*, second edition, Oxford University Press, Oxford, 1998.
- [8] Polterovich L. *The geometry of the group of symplectic diffeomorphisms*. Lectures in Mathematics ETH Zürich Birkhäuser Verlag, Basel, 2001.
- [9] Thèret, David. *Rotation numbers of Hamiltonian isotopies in complex projective spaces*. Duke Math. J. **94**(1998), no. 1, 13-27.
- [10] Weinstein, A. *Cohomology of Symplectomorphism Groups and Critical Values of Hamiltonians*. Math. Zeit., 201, 75-82, 1989.